

What Do Gravitational Lens Time Delays Measure?

C.S. Kochanek

Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138

email: ckochanek@cfa.harvard.edu

ABSTRACT

Gravitational lens time delays depend on the Hubble constant, the observed image positions, and the surface mass density of the lens in the annulus between the images. Simple time delay lenses like PG1115+080, SBS1520+530, B1600+434, PKS1830-211 and HE2149-2745 have $H_0 = A(1 - \langle\kappa\rangle) + B\langle\kappa\rangle(\eta - 1)$ where the two coefficients $A \simeq 90$ km/s Mpc and $B \simeq 10$ km/s Mpc depend on the measured delays and the observed image positions, $\langle\kappa\rangle$ is the mean surface density in the annulus between the images, and there is a small correction from the logarithmic slope $\eta \simeq 2$ of the surface density profile, $\kappa \propto R^{1-\eta}$, in the annulus. These 5 systems are very homogeneous, since for fixed $H_0 = 100h$ km/s Mpc they must have the same surface density, $\langle\kappa\rangle = 1.11 - 1.22h \pm 0.04$, with an upper bound of $\sigma_\kappa < 0.07$ on any dispersion in $\langle\kappa\rangle$ beyond those due to the measurement errors. If the lenses have their expected dark halos, $\langle\kappa\rangle \simeq 0.5$ and $H_0 \simeq 51 \pm 5$ km/s Mpc, while if they have constant mass-to-light ratios, $\langle\kappa\rangle \simeq 0.1-0.2$ and $H_0 \simeq 73 \pm 8$ km/s Mpc. More complicated lenses with multiple components or strong perturbations from nearby clusters, like RXJ0911+0551 and Q0957+561, are easily recognized because they have significantly different coefficients.

Subject headings: cosmology: gravitational lensing; cosmology: Hubble constant; dark matter

1. Introduction

Gravitational lens time delay measurements can determine the Hubble constant H_0 given a model for the gravitational potential of the lens galaxy (Refsdal 1964). With 9 well-measured time delays (see Schechter 2000), gravitational lenses have become a serious alternative to the local distance scale. Kochanek (2002a) modeled the 5 simple, well-observed time delay lenses to find that $H_0 = 48^{+7}_{-4}$ km/s Mpc if the lenses have isothermal mass distributions (flat rotation curves) and that $H_0 = 71 \pm 6$ km/s Mpc if they have constant mass-to-light (M/L) ratios. Intermediate values can be found by adjusting the dark matter halo between these two extreme models, but the results agree with the local estimate of $H_0 = 72 \pm 8$ km/s Mpc by the HST Key Project (Freedman et al. 2001) only if the lenses contain little or no dark matter. Models of other lenses

(e.g. Munoz, Kochanek & Keeton 2001), local stellar dynamical measurements (e.g. Rix et al. 1997, Romanowsky & Kochanek 1999, Gerhard et al. 2001, Treu & Koopmans 2002), weak lensing (e.g. Guzik & Seljak 2002) and X-ray (e.g. Fabbiano 1989, Lowenstein & White 1999) measurements all require mass distributions close to the dark matter limit.

The dependence of the estimate of H_0 from a gravitational lens time delay on the mass distribution of the lens is well known, both from models of particular time delay lenses (e.g. Keeton & Kochanek 1997, Impey et al. 1998, Koopmans & Fassnacht 1999, Lehar et al. 2000, Keeton et al. 2000, Williams & Saha 2000, Winn et al. 2002) and from general analytic principles (Falco et al. 1985, Gorenstein et al. 1988, Refsdal & Surdej 1994, Witt, Mao & Schechter 1995, Witt, Mao & Keeton 2000, Schechter 2000, Rusin 2000, Saha 2000, Zhao & Pronk 2001, Wucknitz 2002, Oguri et al. 2002). The most important trend is that the predicted time delays, or the inferred H_0 , increase as the mass distribution becomes more centrally concentrated. In particular, Falco et al. (1985) showed that adding a constant surface density sheet to the mass distribution has no observable effect other than to rescale the time delay (the mass sheet degeneracy), and Witt et al. (2000) showed that the Hubble constants estimated from lens potentials of the form $\phi \propto R^{3-\eta}$ ($\rho \propto r^{-\eta}$) roughly scale as $H_0 \propto (\eta - 1)/\Delta t$, almost independent of the angular structure of the potential.

Our objective in this paper is to clarify the physical properties of the lenses which control the expected time delays after making full use of the astrometric constraints available for a typical lens. We will show that time delays are controlled by a local property of the lens, the average surface density in the annulus between the lensed images. Reducing the model-dependence of the time delay estimates to such a simple physical property of the lens leads to simple, accurate scaling laws for H_0 , provides model-independent tests for the homogeneity of the time delay lenses, demonstrates that standard parametric models have the necessary degrees of freedom to study degeneracies in estimates of H_0 from time delay measurements, and makes it easy to theoretically predict time delays for standard halo models. In §2 we develop analytic results for lenses in arbitrary spherical density distributions. We first consider circular lens potentials (§2.1) and then examine the effects of adding an external shear (§2.2), the quadrupole of a singular isothermal ellipsoid (§2.3) or a general quadrupole (§2.4). These analytic results show that the time delays are largely determined by the mean surface density $\langle \kappa \rangle = \langle \Sigma \rangle / \Sigma_c$ in the annulus between the images. In §3 we show that the H_0 estimates for most lenses can be reduced to a linear function of $\langle \kappa \rangle$, and that these simple scaling solutions reproduce the full numerical models by Kochanek (2002a). In §4 we discuss the consequences of these results for the homogeneity of the lens population.

2. Analytic Models

We start from the dimensionless time delay (see Schneider, Ehlers & Falco 1992),

$$\tau = \frac{1}{2} (x - u)^2 + \frac{1}{2} (y - v)^2 - \phi(x, y) \quad (1)$$

where x and y are the angular coordinates on the lens plane, u and v are angular coordinates on the source plane, and $\phi(x, y)$ is the lens potential. The center of the lens galaxy is used as the coordinate origin. The true delay is

$$t = \frac{1 + z_l}{H_0} \frac{D_{OL} D_{OS}}{r_H D_{LS}} \tau \equiv \hat{D}_{eff} H_0^{-1} \tau \quad (2)$$

where the D_{ij} are angular diameter distances between the observer, lens and source, z_l is the lens redshift, and H_0 is the Hubble constant.¹ The factor containing the distances and the Hubble radius, $r_H = c/H_0$, depends on the cosmological model, the source redshift z_s and the lens redshift z_l , but not the Hubble constant. We assume a $\Omega_0 = 0.3$ flat universe, but the cosmological dependence of the results is very weak. The lens potential satisfies the Poisson equation $\nabla^2 \phi = 2\kappa$ where $\kappa = \Sigma/\Sigma_c$ is the surface density of the lens in units of the critical surface density $\Sigma_c = c^2 D_{OS}/4\pi G D_{OL} D_{LS}$. The first two terms of the time delay (Eqn. 1) form the geometric delay from the bending of the rays, and the remaining term is the Shapiro delay from passing through the gravitational potential of the lens. We observe images at the solutions of $\nabla \tau = 0$.

We derive the time delay differences for a pair of images in a series of analytic models based on multipole expansions for the lens potential (see Kochanek 1991 and Trotter, Winn & Hewitt 2000 for other applications of multipole methods to gravitational lenses). Our objective is to achieve expressions for the delays which are accurate to $\lesssim 5\%$ when estimating the Hubble constant. Two images at radius R_1 and R_2 have an average radius of $\langle R \rangle = (R_1 + R_2)/2$ and bracket an annulus of width $\Delta R = R_1 - R_2$. For the monopole of the lens, our accuracy goal will require expansions of order $(\Delta R/\langle R \rangle)^3$. The ellipticity of the lens potential generally satisfies $\epsilon < \Delta R/\langle R \rangle$ for two-image lenses and $\epsilon \sim \Delta R/\langle R \rangle$ for four-image lenses. Assuming a standard multipole sequence for a roughly ellipsoidal mass distribution, the quadrupole ($\exp(2i\theta)$) is of order ϵ and can be treated to one lower order of expansion than the monopole, and higher order poles ($\exp(2mi\theta)$) can be neglected since they are of order ϵ^m . We start in §2.1 by estimating the time delays for an arbitrary spherical lens. Next we add a quadrupole potential to the lens: in §2.2 we consider an external (tidal) shear, in §2.3 we consider the quadrupole produced by a singular isothermal ellipsoid, and in §2.4 we consider a quadrupole with an arbitrary ratio between the internal and external contributions to the quadrupole near the images.

¹We note in passing that the distance factor scaling of the time delays is simpler if comoving distances are used rather than angular diameter distances. The extra factor of $1 + z_l$ is absorbed into the distances. Comoving distances, not angular diameter distances, are the natural cosmological distances for gravitational lensing calculations as they almost always lead to simpler analytic expressions.

2.1. Spherical Models

We start with a lens having two images on opposite sides of a spherical lens, $\phi(x, y) = \phi_0(R)$, defined by a monopole potential $\phi_0(R)$. The outer image, at radius R_1 , is at a minimum of the time delay surface and the inner image, at radius R_2 , is at a saddle point, and the two images and the lens lie on a line. Starting from Eqn. (1), the geometric delay between the two images can be written

$$\Delta\tau_{12,geom} = -\frac{1}{2} (R_1^2 - R_2^2) (1 - 2\langle\kappa\rangle) \quad (3)$$

where

$$\langle\kappa\rangle = \frac{2 \int_{R_1}^{R_2} \kappa(u) u du}{R_2^2 - R_1^2} \quad (4)$$

is the average surface density in the annulus bounded by the images in units of the critical surface density. Only the derivative of the potential is needed to determine the geometric delay, and the mean surface density enters because the radial derivatives ϕ'_0 of the monopole at the positions of the two images are related by

$$R_2\phi'_0(R_2) = R_1\phi'_0(R_1) + \langle\kappa\rangle(R_2^2 - R_1^2) \quad (5)$$

independent of the density distribution in the annulus. For an isothermal lens $\langle\kappa\rangle = 1/2$ and the geometric contribution to the delay is identically zero for all images. The Shapiro delay difference is more complicated,

$$\Delta\tau_{12,Shap} = \frac{1}{2}\langle\kappa\rangle (R_2^2 - R_1^2) + R_1 R_2 (1 - \langle\kappa\rangle) \ln \frac{R_2}{R_1} - 2 \int_{R_1}^{R_2} u du (\kappa(u) - \langle\kappa\rangle) \ln \frac{u}{R_2}, \quad (6)$$

because the Shapiro delay depends on the surface density distribution in the annulus as well as the mean surface density. The first term, which depends only on $\langle\kappa\rangle$, is the most important, as the contribution from the integral is smaller by $(\Delta R/\langle R\rangle)^2$. Combining the terms, the time delay between the images,

$$\Delta\tau_{12} = \tau_1 - \tau_2 = (1 - \langle\kappa\rangle) \left[-\frac{1}{2} (R_1^2 - R_2^2) + R_1 R_2 \ln \frac{R_2}{R_1} \right] - 2 \int_{R_1}^{R_2} u du (\kappa(u) - \langle\kappa\rangle) \ln \frac{u}{R_2}, \quad (7)$$

depends only on the image positions and the surface density of the lens in the annulus between them (Gorenstein et al. 1988). The first term is the same as the time delay of a point mass lens combined with a uniform sheet of density $\langle\kappa\rangle$.

We can capture the essential elements of Eqn. (7) without the complexity by assuming that the surface density in the annulus between the images is a power law $\kappa = \kappa_1(R/R_1)^{1-\eta}$ and then expanding the time delay as a series in the ratio of the thickness of the annulus $\Delta R = R_2 - R_1$ to its average radius $\langle R \rangle = (R_1 + R_2)/2$ to find that

$$\Delta\tau_{12} = - (R_1^2 - R_2^2) \left[(1 - \langle\kappa\rangle) - \frac{1 - \eta\langle\kappa\rangle}{12} \left(\frac{\Delta R}{\langle R \rangle} \right)^2 + O \left(\left(\frac{\Delta R}{\langle R \rangle} \right)^4 \right) \right]. \quad (8)$$

The first term is the exact delay for a singular isothermal sphere (SIS, $\eta = 2$, $\langle\kappa\rangle = 1/2$), and the series converges rapidly even for large values of $\Delta R/\langle R\rangle$. For example, the first term approximates the time delay of a point mass lens with only a 9% error even as $\Delta R/\langle R\rangle \rightarrow 1$. The second order correction adjusts for the small changes in the critical radius (inside which the average surface density is unity) with the shape of the density profile. It comes only from the Shapiro delay (Eqn. 7) and can be decomposed ($1 - \eta\langle\kappa\rangle = (1 - \langle\kappa\rangle) + \langle\kappa\rangle(\eta - 1)$) into a contribution depending only on the average surface density and a contribution which depends on its distribution. The dependence on the logarithmic slope η arises only from the integral term of the Shapiro delay. The scaling observed by Witt et al. (2000) for potentials of the form $\phi \propto R^{3-\eta}$ arises because $1 - \langle\kappa\rangle = (\eta - 1)/2$ near the critical line of the lens.

2.2. A Spherical Lens in an External Shear

Real lenses are not spherical, so we next consider a range of models with a quadrupole as well as a monopole. The monopole depends only on the total mass inside the Einstein ring and the surface density in the annulus. The quadrupole depends on the total internal quadrupole (due to the quadrupole moment of the mass inside the Einstein ring), the total external quadrupole (due to the quadrupole moment of the mass outside the ring) and the quadrupole component from the surface density inside the annulus. Because the ellipticity of the potential is small ($\lesssim \Delta R/\langle R\rangle$) we can treat the quadrupole to a lower order of expansion than the monopole and ignore the quadrupole component of the surface density in the annulus. For our analytic results, but not for the numerical results in §3, we assume that the internal and external quadrupoles are aligned. Thus, we need three numbers, two amplitudes and an orientation, to define the quadrupole. Higher order angular structures (octopole etc.) will be smaller than the quadrupole by one or more powers of the ellipticity of the potential, and can be neglected.

We start with two familiar examples of quadrupoles. We first consider an external (tidal) shear, where the quadrupole is generated entirely by material outside the region with the images. In §2.3 we discuss the quadrupole produced by a singular isothermal ellipsoid. Then in §2.4 we discuss the general aligned quadrupole. Ideally, we would present simple results to the same order ($(\Delta R/\langle R\rangle)^3$) as we did for the monopole. However, our objective is to illustrate the effects of quadrupole structures on time delays, so we will not reproduce terms which are grotesquely complicated simply to reach an arbitrary order of expansion.

The quadrupole potential associated with an external shear is $\phi_2 = (\gamma_{ext}/2)R^2 \cos 2(\theta - \theta_{ext})$, where γ_{ext} is the amplitude of the external shear and θ_{ext} is its orientation. As in §2.1, the monopole is defined by the mass inside the Einstein ring, the average surface density $\langle\kappa\rangle$ in the annulus and the logarithmic derivative of the surface density η . We model two images located at $(R_1 \cos \theta_1, R_1 \sin \theta_1)$ and $(R_2 \cos \theta_2, R_2 \sin \theta_2)$ from the lens center. Using the positions of two images we solve for the mass inside the Einstein ring and the amplitude of the external shear. This results in expressions which depend on the surface density in the annulus ($\langle\kappa\rangle$, η) and the

orientation of the shear (θ_{ext}). The shear axis seems the best choice for an independent variable in our analytic models because its value is little affected by changes in the monopole or quadrupole structure of the lens (see Kochanek 1991). If we again expand the delay as a series in $\Delta R/\langle R \rangle$,

$$\Delta\tau_{12} = -\left(R_1^2 - R_2^2\right) \left[T_0 + \frac{\Delta R}{\langle R \rangle} T_1 + \left(\frac{\Delta R}{\langle R \rangle}\right)^2 T_2 \cdots \right] \quad (9)$$

then the leading term,

$$T_0 = \left(1 - \langle \kappa \rangle\right) \sin^2 \left(\frac{\Delta\theta_{12}}{2}\right) \longrightarrow \left(1 - \langle \kappa \rangle\right) \quad \text{as} \quad \Delta\theta_{12} \rightarrow \pi, \quad (10)$$

changes from the result for a spherical system only in adding a dependence on the angle between the images $\Delta\theta_{12} = \theta_1 - \theta_2$. The limit as $\Delta\theta_{12} \rightarrow \pi$ shows that we recover the result for a circular lens and images collinear with the lens (Eqn. 8). The first order correction,

$$T_1 = -\frac{1}{2} \left(1 - \langle \kappa \rangle\right) \sin(\Delta\theta_{12}) \cot(\theta_1 + \theta_2 - 2\theta_{ext}) \longrightarrow 0 \quad \text{as} \quad \Delta\theta_{12} \rightarrow \pi, \quad (11)$$

depends on the shear axis θ_{ext} , but the second order correction,

$$T_2 = \frac{1}{12} \left[-\left(1 - \langle \kappa \rangle\right) (4 + 3 \cos \Delta\theta_{12}) + \langle \kappa \rangle (\eta - 1) \right] \longrightarrow -\frac{1 - \eta \langle \kappa \rangle}{12} \quad \text{as} \quad \Delta\theta_{12} \rightarrow \pi, \quad (12)$$

depends only on the surface density in the annulus. Note that the higher order terms return to the spherical case for images collinear with the lens. The amplitude of the shear is

$$\gamma_{ext} = -\frac{(1 - \langle \kappa \rangle) (R_1^2 - R_2^2)}{(R_1^2 + R_2^2 - 2R_1 R_2 \cos \Delta\theta_{12}) \sin(\theta_1 + \theta_2 - 2\theta_{ext})} \longrightarrow 0 \quad \text{as} \quad \Delta\theta_{12} \rightarrow \pi, \quad (13)$$

so the singularity in T_1 when $\theta_1 + \theta_2 - 2\theta_{ext} \rightarrow 0$ is due to a singularity in the shear that will not arise in practice since θ_{ext} must be set using the parameters of a reasonable macro model. Also note that the amplitude of the shear scales as $1 - \langle \kappa \rangle$, so that lenses with lower surface densities require larger shears.

The dependence of the delay on the structure of the surface density in the annulus is identical to that for the spherical lens, with the integral in the Shapiro delay (Eqn. 6) adding a term $(\eta - 1)\langle \kappa \rangle/12$ to the T_2 coefficient. All other terms in the time delay depend only on the derivatives of the monopole and are functions only of $\langle \kappa \rangle$ even for an arbitrary monopole surface density. The term vanishes for $\eta = 1$ when $\kappa = \kappa_1(R/R_1)^{1-\eta}$ becomes a constant surface density.

2.3. A Spherical Lens in an SIE Quadrupole

Unlike our result for an external shear, where the time delay depends on the angle between the images even at the lowest order of the expansion, Witt et al. (2000) found that the time delays for potentials of the form $\phi \propto R\mathcal{F}(\theta)$ (which includes the standard singular isotherma ellipsoid or

SIE potential) depend only on the image radii. These models have a quadrupole potential of the form $\phi_2 = \epsilon R \langle R \rangle \cos 2(\theta - \theta_\epsilon)$. The leading term in the time delay,

$$T_0 = 1 - \langle \kappa \rangle \quad (14)$$

is independent of the angle, just as for the Witt et al. (2000) potential. The second term is a correction for the effects of the ellipticity,

$$T_1 = -\frac{1}{8} (1 - \langle \kappa \rangle) [5 \cos(\Delta\theta_{12}/2) + \cos(3\Delta\theta_{12}/2)] \cot(\theta_1 + \theta_2 - 2\theta_{ext}) \csc^3(\Delta\theta_{12}/2) \quad (15)$$

The next term is too complicated to present in full, but it has the structure

$$T_2 = (1 - \langle \kappa \rangle) g(\theta_1, \theta_2, \theta_\epsilon) + \frac{1}{12} (\eta - 1) \langle \kappa \rangle. \quad (16)$$

The first factor depends on a very complicated angular function, and the second factor is the correction to the Shapiro delay from the logarithmic slope η of the monopole surface density. To lowest order, the ellipticity of the potential is

$$\epsilon \simeq -\frac{1 - \langle \kappa \rangle}{4} \frac{\Delta R}{\langle R \rangle} \sin \Delta\theta_{12} \csc^4(\Delta\theta_{12}/2) \csc(\theta_1 + \theta_2 - 2\theta_\epsilon). \quad (17)$$

We again find that the ellipticity scales as $1 - \langle \kappa \rangle$ and that the singularities in T_1 correspond to singularities in ϵ .

2.4. A Spherical Lens in a General Quadrupole

The quadrupoles of an external shear and an SIE differ in the balance between internal and external quadrupoles. Since they also produce different angular dependences for the leading term in the time delay, it suggests that the angular dependence of the time delay is determined by the relative strength of the two shear components. We can test this by determining the leading term in the time delay for a quadrupole which is an arbitrary sum of an internal quadrupole of amplitude γ_{int} and external quadrupole of amplitude γ_{ext} ,

$$\phi_2 = \frac{1}{2} \left(\gamma_{ext} R^2 + \gamma_{int} \frac{\langle R \rangle^4}{R^2} \right) \cos 2(\theta - \theta_\gamma). \quad (18)$$

If we use the image positions to determine the total shear, $\Gamma = \gamma_{int} + \gamma_{ext}$, as a function of the shear angle θ_γ and the ratio $f_{int} = \gamma_{int}/\Gamma$, we find that the leading term,

$$T_0 = - (1 - \langle \kappa \rangle) \frac{\sin^2(\Delta\theta_{12}/2)}{1 - 4f_{int} \cos^2(\Delta\theta_{12}/2)}, \quad (19)$$

has the standard scaling with $1 - \langle \kappa \rangle$ and an angular dependence determined by the balance between the internal and external quadrupoles. For an external shear ($f_{int} = 0$) the denominator is

unity, and for the quadrupole of an SIE ($f_{int} = 1/4$ for $\kappa_2 \propto R^{-1}$) the numerator and denominator cancel to make T_0 independent of $\Delta\theta_{12}$ just as in the Witt et al. (2000) results. For $f_{int} > 1/4$, where the internal shear begins to dominate, there are angles where the expansion fails. The higher order terms are too complex to be worth presenting for general configurations. However, for two-image lenses where the images lie on opposite sides of the lens, $\theta_2 = \theta_1 + \pi + \delta\theta$, with a small angular offset defined by $\xi = \delta\theta\langle R\rangle/\Delta R \lesssim 1$, the delay depends little on the structure of the quadrupole. The expansion terms,

$$T_0 = 1 - \langle\kappa\rangle, \quad T_1 = 0 \quad \text{and} \quad T_2 = -\frac{1}{12}(1 - \eta)\langle\kappa\rangle + \frac{1}{4}\xi \left(1 - \langle\kappa\rangle\right) [2 \cos 2\theta_\gamma + \xi(4f_{int} - 1)] \quad (20)$$

are nearly identical to those for a spherical lens (Eqn. 8) because the quadrupole structure enters only at second order.

3. A Simple Semi-Analytic Model

These analytic results suggest a simple semi-analytic model for understanding the time delays observed in gravitational lenses. It cannot be a completely analytic model because we require a numerical solution using all the lensed images to determine the quadrupole structure of the four-image lenses. We continue to model the monopole by the mass inside the average image radius and a power law, $\kappa(R) \propto R^{1-\eta}$, for the surface density between the inner and outer images parameterized by the mean surface density $\langle\kappa\rangle$ and the logarithmic slope η . The resulting expression for the monopole deflection,

$$\phi'_0(R) = b_0 \frac{\langle R \rangle}{R} + \frac{2}{R} \int_{\langle R \rangle}^R \kappa(R) R dR, \quad (21)$$

where $b_0 \sim \langle R \rangle$ determines the mass inside $\langle R \rangle$ is very similar to an early model of Q0957+561 by Borgeest & Refsdal (1984) which combined a point mass and an isothermal sphere ($\eta = 2$). Since many of the lenses have $\Delta R/\langle R \rangle \sim 0.5$ we expanded the monopole to first order in the deviation of the density exponent from isothermal, $\delta\eta = \eta - 2$, rather than $\Delta R/\langle R \rangle$. This leads to results that are exact for $\eta = 2$ independent of the width of the annulus, and are reasonably accurate for $1 \lesssim \eta \lesssim 3$. We combined the monopole with a general quadrupole

$$\phi_2(R, \theta) = \frac{1}{2}\gamma_{ext}R^2 \cos 2(\theta - \theta_{ext}) + \frac{1}{2}\gamma_{int}\frac{\langle R \rangle^4}{R^2} \cos 2(\theta - \theta_{int}) \quad (22)$$

where the internal and external components have amplitudes and orientations of $(\gamma_{int}, \theta_{int})$ and $(\gamma_{ext}, \theta_{ext})$ respectively. For each lens we determined the values of b_0 and the shear components which minimized the differences in the projected source positions, leaving expressions depending only on the monopole structure in the annulus ($\langle\kappa\rangle$ and η). For the current time delay lenses, we cannot determine $\langle\kappa\rangle$ or η from the available constraints. By doing the fits on the source plane, the solutions can be found rapidly using *mathematica* programs. We did not fit the image flux

ratios. For the two-image lenses, where there are too few constraints to determine all 5 variables, we stabilized the solutions by finding the minimum shear solutions.² We experimented with the structure of the shear terms (and the shear prior for the two-image lenses) to ensure that the results reported for H_0 are not affected by the assumed quadrupole structure. For the two-image lenses the results had no significant dependence on the quadrupole structure (as expected from §2.4), and for the four-image lenses the quadrupole structure was well-determined by the image positions. Errors were estimated by Monte Carlo simulations and for H_0 they are dominated by the errors in the time delay measurements.

Once we have solved for the model parameters required to match the image positions, we derive the time delays between images as a function of the remaining parameters $\langle\kappa\rangle$ and η . Based on the analytic and numerical results, we can express the results as simple scaling relations for the Hubble constant of the form

$$H_0 \simeq A(1 - \langle\kappa\rangle) + B\langle\kappa\rangle(\eta - 1) + C \quad (23)$$

where the three coefficients (A , B and C) have the units of km/s Mpc and scale inversely with the measured time delays. We have isolated the terms more physically here than in the expansions of §2, isolating the dependence of the logarithmic slope η in the B coefficient, but the relationship between the coefficients and the expansion can be worked out for each model. For example, the T_0 term defined in Eqn. (9) contributes $-(R_1^2 - R_2^2)T_0\hat{D}_{eff}/\Delta t_{12}$ to the A coefficient, where Δt_{12} is the measured delay and $\hat{D}_{eff}$ is the effective distance defined in Eqn. (2). In the analytic models $C \equiv 0$. For simple lenses the Hubble constant $H_0 \simeq A$ for very centrally concentrated lenses ($\langle\kappa\rangle \rightarrow 0$), and $H_0 \simeq (A + B)/2$ for an isothermal model ($\langle\kappa\rangle = 1/2$, $\eta = 2$). We order the images so that the time delay is positive and $A > 0$. The effects of an additional mass sheet of density κ_{ext} can be easily added to the models. The adjustments depend on whether we regard $\langle\kappa\rangle$ as a fixed number or a fit parameter whose value is also changed by the addition of the extra convergence. If $\langle\kappa\rangle$ is a fixed number, then the estimate of the Hubble constant changes to

$$H_0 \simeq (1 - \kappa_{ext}) \left[A(1 - \langle\kappa\rangle) + C \right] + B\langle\kappa\rangle(\eta - 1). \quad (24)$$

In most cases, however, we start with a lens whose mass and surface density are scaled to fully explain the image splitting. With the addition of an external convergence, the annular surface density $\langle\kappa\rangle_0$ of this reference model is reduced to $\langle\kappa\rangle = (1 - \kappa_{ext})\langle\kappa\rangle_0$ because the extra convergence reduces the mass of the primary lens. Thus, if we specify $\langle\kappa\rangle_0$, the estimate of the Hubble constant with the addition of an external convergence is

$$H_0 \simeq (1 - \kappa_{ext}) \left[A(1 - \langle\kappa\rangle_0(1 - \kappa_{ext})) + B\langle\kappa\rangle_0(\eta - 1) + C \right] \quad (25)$$

and it has the familiar $H_0 \propto (1 - \kappa_{ext})$ scaling of the mass-sheet degeneracy (Falco et al. 1985).

²For two-image lenses we added on weak prior on the shear amplitudes to the χ^2 for the mapping of the images back to a common source. The prior was defined by $\mathbf{\Gamma} \cdot \mathbf{\Gamma}\gamma_0^{-2}$ where $\mathbf{\Gamma} = (\gamma_{int} \cos 2\theta_{int}, \gamma_{int} \sin 2\theta_{int}, \gamma_{ext} \cos 2\theta_{ext}, \gamma_{ext} \sin 2\theta_{ext})$ is the vector of Cartesian shear components and we set $\gamma_0 = 0.05$.

Table 1. Hubble Constant Models

Lens	Images	Delay (days)	$R_e/\langle R \rangle$	$\frac{\Delta R}{2\langle R \rangle}$	A_0 [H_0]	A [H_0]	B [H_0]	C [H_0]
RXJ0911+0551	AD	146 ± 4	0.55	0.58	272 ± 14	82 ± 5	69 ± 4	-9 ± 1
	BD			0.54	280 ± 14	81 ± 5	69 ± 5	-9 ± 1
	CD			0.60	262 ± 14	82 ± 6	68 ± 5	-9 ± 1
Q0957+561		417 ± 3	0.63	0.67	249 ± 2	221 ± 2	51 ± 1	-7 ± 0
PG1115+080	A ₁ B	12 ± 2	0.41	0.10	81 ± 17	70 ± 14	0 ± 0	1 ± 0
	A ₁ C	13 ± 2		0.10	126 ± 22	112 ± 20	9 ± 2	0 ± 0
	BC	25 ± 2		0.19	97 ± 8	93 ± 8	5 ± 0	0 ± 0
SBS1520+530		129 ± 3	0.51	0.52	98 ± 5	92 ± 5	10 ± 1	-1 ± 0
B1600+434		51 ± 2	0.94	0.64	119 ± 14	104 ± 10	21 ± 9	-3 ± 2
PKS1830-211		26 ± 4		0.19	85 ± 15	88 ± 16	1 ± 0	0 ± 0
HE2149-2745		103 ± 12	0.55	0.59	95 ± 12	84 ± 11	14 ± 2	-1 ± 0

Note. — For each lens and image pair we give the measured time delay in days, followed by the coefficients which determine the Hubble constant given the surface density model. We first give the single coefficient A_0 needed for the lowest order expansion $H_0 = A_0(1 - \langle \kappa \rangle)$ for an arbitrary monopole in an external shear, and then the three coefficients A , B and C used in our standard expansion (Eqn. 23) for an arbitrary monopole and quadropole. The formal errors include the uncertainties in both the astrometry and time delays but are dominated by the time delays. This means that the uncertainties are almost perfectly correlated! We broaden the time delay errors to a minimum error of 5% to account for systematic uncertainties such as convergence fluctuations from large scale structure. The ratio $R_e/\langle R \rangle$ is the ratio of the (monopole) half-light radius to the average image radius, and the ratio $\Delta R/2\langle R \rangle$ is an estimate of the thickness of the annulus.

Table 2. Comparisons to Full Numerical Models

Lens	Image	Full Model H_0		$\langle\kappa\rangle_0/\kappa_{ext}$		Lowest Order H_0		Standard H_0	
		DM	M/L	DM	M/L	DM	M/L	DM	M/L
RXJ0911+0551	AD	49 ± 4	67 ± 4	0.50/0.24	0.11/0.31	129 ± 7	174 ± 9	58 ± 3	...
	BD			0.50/0.24	0.10/0.31	132 ± 7	180 ± 9	58 ± 3	...
	CD			0.50/0.24	0.12/0.31	123 ± 7	166 ± 9	58 ± 3	...
PG1115+080	A ₁ B	46 ± 4	69 ± 5	0.50/0.09	0.18/0.16	40 ± 8	58 ± 12	35 ± 7	51 ± 10
	A ₁ C			0.50/0.09	0.11/0.16	62 ± 11	96 ± 17	60 ± 10	87 ± 15
	BC			0.50/0.09	0.15/0.16	48 ± 4	71 ± 6	48 ± 4	69 ± 6
SBS1520+530		53 ± 6	73 ± 5	0.50/0.00	0.20/0.00	49 ± 3	79 ± 4	51 ± 3	77 ± 4
B1600+434		59 ± 6	71 ± 9	0.50/0.05	0.43/0.18	60 ± 7	63 ± 7	59 ± 7	68 ± 9
HE2149–2745		47 ± 8	66 ± 8	0.50/0.00	0.22/0.00	48 ± 6	74 ± 9	48 ± 6	70 ± 9

Note. — Hubble constant estimates for dark matter (DM) and constant mass-to-light ratio (M/L) models of the density distribution. All estimates of the Hubble constant are in units of km/s Mpc. The estimates from full numerical models are the dark matter and constant M/L models from Kochanek (2002a). The lowest order H_0 estimates use the lowest order expansion $H_0 = A_0(1 - \langle\kappa\rangle)$, while the standard H_0 estimates use Eqn. (23). The $\langle\kappa\rangle_0/\kappa_{ext}$ columns give the mean surface density $\langle\kappa\rangle_0$ of an unperturbed spherical model and the surface density due to external perturbers κ_{ext} in the full numerical models. For the isothermal models we used $\eta = 2$ while for the constant M/L models we used $\eta = 3$.

We computed these simple models for 7 of the 9 time delay lenses, excluding only B0218+357 (where the lens position is poorly known, Lehar et al. 2000) and B1608+656 (where the lens consists of two interacting galaxies, Koopmans & Fassnacht 1999). We consider RXJ0911+0551 (Hjorth et al. 2002), Q0957+561 (Schild & Thomson 1997, Kundic et al. 1997, Bernstein et al. 1997, Keeton et al. 2000), PG1115+080 (Schechter et al. 1997, Barkana 1997, Impey et al. 1998), SBS1520+530 (Burud 2002b, Faure et al. 2002), B1600+434 (Burud et al. 2000, Koopmans et al. 2000), PKS1830–211 (Lovell et al. 1998, Winn et al. 2002, Courbin et al. (2002)) and HE2149–2745 (Burud et al. 2002). The system parameters are generally derived from the CfA/Arizona Space Telescope Lens Survey (CASTLES, Falco et al. 2001) images of the systems.

Of these 7 systems, we expect our simple models to have difficulty with Q0957+561 and RXJ0911+0551. Q0957+561 is near the center of a cluster (Keeton et al. 2000, Chartas et al. 2002) where an external shear will poorly describe the effect of the cluster on the image positions. RXJ0911+0551 is on the outskirts of a cluster (see Morgan et al. 2001) and the primary lens also has a small satellite. The external shear approximation is a poor one for the perturbations from the RXJ0911+0551 cluster. It is at best a marginal approximation for $\langle\kappa\rangle \sim 1/2$ and it fails completely for $\langle\kappa\rangle \sim 0$ (recall that $\gamma \propto (1 - \langle\kappa\rangle)$, see Eqns. 13 and 17). We included an SIS satellite in our model of RXJ0911+0551, so it will match the Kochanek (2002a) dark matter models, but it is difficult to treat the two potentials self-consistently within our approximation. We also note that our estimates for PKS1830–211 are invalid if we adopt the Courbin et al. (2002) multiple-lens interpretation of the system rather than the Winn et al. (2002) single-lens interpretation.

We now check our approximations against full numerical models for RXJ0911+0551, PG1115+080, SBS1520+530, B1600+434 and HE2149–2745 by Kochanek (2002a). We first checked the dependence of the results on the assumed quadrupole structure by fitting models with either a pure external shear, an SIE quadrupole or a pure internal shear. For the two-image lenses, the H_0 estimates are essentially independent of the quadrupole structure, as expected from the analysis in §2.4. The largest fractional changes in H_0 are approximately 3%. This is fortunate because the observational constraints on the two-image lenses cannot distinguish between the quadrupole structures. When we allow a general quadrupole, which matches realistic models where there are contributions to the angular structure from both tidal shears and the ellipticity of the lens galaxy (Keeton, Kochanek & Seljak 1997), all the doubles can be fit perfectly and have H_0 estimates consistent with the three constrained shear models.

For the two four-image lenses, RXJ0911+0551 and PG1115+080, the quadrupole structure is enormously important because the functional dependence of the delay on the angle between the images is so strong once the images do not lie on opposite sides of the lens. For example, for an SIS monopole ($\langle\kappa\rangle = 1/2$, $\eta = 2$), the H_0 estimates from the BC image pair in PG1115+080 are 47 and 67 km/s Mpc for an external shear and an SIE quadrupole respectively. The ratio of the values matches that expected from Eqn. (19). Because they are four-image lenses, however, they also strongly distinguish between quadrupole structures, with both lenses being far more consistent with an external shear than either a SIE quadrupole or an internal shear. Time delay ratios

are also controlled by the quadrupole structure, and the measured delay ratios in PG1115+080 are also more consistent with an external shear than either an SIE quadrupole or an internal shear. The delay ratio seemed to be model independent in our studies of PG1115+080 (Keeton & Kochanek 1997, Impey et al. 1998) because the models focused on changes in the monopole while always optimizing the quadrupole structure. Since the best fit quadrupole was close to an external shear no matter what we used for the monopole structure, the time delay ratios showed almost no model dependence. For models with a general quadrupole, we find that $f_{int} \simeq 0.02$ and 0.01 for RXJ0911+0551 and PG1115+080 respectively. While very close to a pure external shear ($f_{int} = 0$), the general quadrupole significantly improves the fit to the image positions.

We adopt the general quadrupole models as our standard. For the two-image lenses the choice has negligible effect on the estimates of H_0 , and for the four-image systems it is required to provide an acceptable fit to the image positions. Table 1 presents the coefficients A , B , and C (Eqn. 23) as well as the coefficient A_0 , with $H_0 = A_0(1 - \langle\kappa\rangle)$, we would obtain using the lowest order term for an external shear model (T_0 from Eqn. 10). The uncertainties in the coefficients are almost entirely due to the time delay estimates, although they do include the astrometric uncertainties in the image and lens positions. These become important only for B1600+434 where the lens position is relatively uncertain due to the dust lane in the galaxy (Koopmans, de Bruyn & Jackson 1998, Maller et al. 2000). The dominant role of the time delay uncertainties means that the uncertainties in the coefficients are almost perfectly correlated. With the exception of the two lenses in clusters, RXJ0911+0551 and Q0957+561, the lowest order coefficient A_0 matches the higher order estimate A to approximately 10%, the B coefficient is needed to achieve 10% accuracy, and the C coefficient is negligible compared to other sources of uncertainty. The amplitude of B is largest for the lenses with thicker annuli (larger $\Delta R/\langle R \rangle$). In fact, the coefficients clearly divide the sample into a set of five very similar, relatively isolated lenses (PG1115+080, SBS1520+530, B1600+434, PKS1830–211, and HE2149–2745) and the two lenses in clusters (RXJ0911+0551 and Q0957+561).

The real question, however, is whether these simple semi-analytic representations can reproduce the results of full numerical models. Table 2 compares the semi-analytic estimates to the full numerical lens models from Kochanek (2002a) for either a dark matter dominated, SIE lens model ($\langle\kappa\rangle_0 = 1/2$, $\eta = 2$) or a constant M/L model where $\langle\kappa\rangle_0 \sim 0.1$ – 0.2 for most lenses. We defined $\langle\kappa\rangle_0$ for the constant M/L models by the average density in the annulus for a spherical model using the intermediate axis effective radius (the geometric mean of the major and minor axes) for the scale length in a de Vaucouleurs profile. For a lens with surface density profile $\Sigma(R)$, the critical surface density profile is simply

$$\kappa(R) = \frac{R_c^2 \Sigma(R)}{2 \int_0^{R_c} R \Sigma(R) dR} \quad (26)$$

where R_c is the average critical radius of the lens. It depends only on the shape of the profile because for a spherical lens the mean density inside the critical radius is equal to the critical density so the image geometry determines the mass scale. We also include the effects of the

external convergences κ_{ext} from the numerical models. We do not include the constant M/L semi-analytic estimates for RXJ0911+0551 where the treatment of the nearby cluster as an external shear becomes untenable.

First, note that the estimates for each image pair in the four-image lenses RXJ0911+0551 and PG1115+080 are mutually consistent. This holds for the A₂B and A₂C pairs of PG1115+080 as well. Second, for the simple lenses, the lowest order and standard approximations are nearly identical for the dark matter (high $\langle\kappa\rangle$) models, but the lowest order approximation tends to give higher estimates for H_0 for the constant M/L (low $\langle\kappa\rangle$) models. For the more complicated lens RXJ0911+0551, the differences are enormous. Finally, the approximate solutions are an excellent match to the full numerical models using ellipsoidal lenses in external shear fields. Using the AD image pair from RXJ0911+0551 and the BC image pair from PG1115+080, the average difference between the approximate and numerical dark matter models is $\Delta H_0 = 2 \pm 4$ km/s Mpc with almost all the differences due to the most complex system, RXJ0911+0551. The approximate model for RXJ0911+0551 completely fails in the constant M/L limit (see above), but if we compare the remaining four lenses the difference is $\Delta H_0 = 2 \pm 1$ km/s Mpc. For comparison, the average differences for the lowest order approximation are $\Delta H_0 = 0 \pm 2$ km/s Mpc and $\Delta H_0 = 2 \pm 6$ km/s Mpc respectively, but we have to drop RXJ0911+0551 for the dark matter models as well as the constant M/L models. Some of these residuals may be due to the ambiguities in defining $\langle\kappa\rangle$ for ellipsoidal lens galaxies. We conclude that our approximate solutions generally provide estimates for H_0 to better than 10% individually and to better than 5% as an ensemble.

4. Discussion

Gravitational lens time delays are determined by the Hubble constant, the positions of the lensed images, and the surface density in the annulus bounded by the images. The average surface density $\langle\kappa\rangle$ in the annulus is more important than its distribution. The relationship between time delays and the local surface density is exact for circular lenses (Gorenstein et al. 1988). While it is not exactly true for non-circular lenses, it is true in practice. In two-image lenses, where the images lie on opposite sides of the lens galaxy, the delays are insensitive to the angular structure of the lens. In four-image lenses the delays are very sensitive to the quadrupole structure of the potential, but the image positions tightly constrain the quadrupole and leave the surface density as the only important variable. Our local interpretation of time delays agrees with earlier understandings based on global models and degeneracies, but better isolates the relevant physical properties of lens models. It also allows us to find simple scaling solutions for the Hubble constant which accurately reproduce the results of full numerical models. These scaling solutions can be used to classify time delay lenses, since their structure differs for simple lenses and lenses strongly perturbed by clusters, and to quickly estimate the Hubble constant predicted for any assumed radial mass distribution.

We can also use the scaling solutions to study the homogeneity of the lens population.

Because H_0 must be a universal constant, we can use the requirement that the lens all produce the same Hubble constant to estimate the surface density differences between the lenses. If we assume a universal density slope η , then the five simple lenses (PG1115+080, SBS1520+530, B1600+434, PKS1830–211 and HE2149–2745) are consistent with a common value for the average surface density of

$$\langle \kappa \rangle \simeq 1 - 1.07h + 0.14(\eta - 1)(1 - h) \pm 0.04, \quad (27)$$

where $H = 100h$ km/s Mpc. Formally, only limited ranges for the Hubble constant are consistent with a universal surface density, with $h < 0.7$, $h = 0.5 \pm 0.6$ and $h = 0.9 \pm 0.3$ for $\eta = 1$ (constant surface density), 2 (isothermal) and 3 respectively. An alternative way to characterize the similarities is use one lens as a reference and then estimate the surface density differences between the reference lens and the other systems. For example, if we assume $\eta = 2$ and use PG1115+080 as the reference system, we find that $\langle \kappa \rangle = 1.05 - 1.13h \pm 0.04$ for PG1115+080. The surface density differences are $\Delta\kappa = 0.07 - 0.09h \pm 0.06$, $0.19 - 0.07h \pm 0.11$, $0.00 - 0.00h \pm 0.19$ and $0.14 - 0.28h \pm 0.15$ for SBS1520+530, B1600+434, PKS1830–211 and HE2149–2745 respectively. The images in PG1115+080, SBS1520+530 and HE2149–2745 all lie at approximately the same radius relative to the lens galaxy ($R_e/\langle R \rangle \simeq 0.5$) while the images in B1600+434 are close to the half-light radius ($R_e/\langle R \rangle \simeq 1$). Reasonable slope differences between the lenses can change the surface density differences by $0.00\text{--}0.27(1 - h)\Delta\eta$ depending on the lens. At least for constant η , the upper bound on the existence of any scatter in the surface density beyond that implied by the measurement errors is $\sigma_\kappa \lesssim 0.07$, so these five simple lenses have very homogeneous intrinsic properties.

Our analysis is not well suited to the two cluster lenses, RXJ0911+0551 and Q0957+561, because our use of an external shear to represent perturbations is breaking down. If, however, we assume that the true surface density of the primary lens is the same as for the other lenses, we can estimate the external convergence κ_{ext} required to make the time delay estimates for the cluster lenses agree with the other 5 systems. We find that $\kappa_{ext,0911} \simeq 1.11 - 1.43h \pm 0.10$ and $\kappa_{ext,0957} \simeq 0.99 - 0.45h \pm 0.01$. While these results should not be taken at face value because of the problems with our approximation for lenses which are too close to the central regions of clusters, the pattern of the solutions seems reasonable. The RXJ0911+0551 lens lies $200h^{-1}$ kpc from the cluster center, while the Q0957+561 lens lies only $30h^{-1}$ kpc from the cluster center so we would expect larger surface densities in Q0957+561 than in RXJ0911+0551. Presumably similar analyses with more realistic models for the cluster would improve matters.

Our semi-analytic models for the four simple lenses with good photometry, PG1115+080, SBS1520+530, B1600+434 and HE2149–2745, almost exactly reproduce the H_0 estimates based on full numerical models by Kochanek (2002a). We find $H_0 \simeq 51 \pm 5$ km/s Mpc if $\kappa = 1/2$ and $\eta = 2$ as expected in dark matter dominated lenses and $H_0 \simeq 73 \pm 8$ km/s Mpc if they have constant mass-to-light ratios. The dependence of the time delays on the radial and angular structure of the lens shows that standard parametric models, which can adjust both the radial mass distribution and the quadrupole structure of the lens, encompass the physical properties

needed to study the dependence of Hubble constant estimates on the mass distribution of the lens. Most degrees of freedom in non-parametric models (e.g. Williams & Saha 2000) are not important for time delay estimates in simple lenses. Since the parametric models have the advantage of corresponding to physical models of galaxies, while most realizations of the non-parametric models do not (see Munoz et al. 2001), the use of non-parametric approaches is probably better suited to very complicated systems like B1608+656 where the lens consists of two interacting galaxies (Koopmans & Fassnacht 1999) and it is unclear how to properly parameterize the system. It is also easy to make theoretical estimates for the time delays expected for standard cold dark matter (CDM) halo models, since we need only estimate the surface density $\langle\kappa\rangle$ of the halo. We examine this problem in Kochanek (2002b).

Acknowledgments. CSK thanks D. Rusin, P. Schechter, J. Winn and S. Wyithe for discussions and comments. CSK is supported by the Smithsonian Institution and NASA ATP grant NAG5-9265.

REFERENCES

- Barkana, R., 1997, *ApJ*, 489, 21
- Bernstein, G., Fischer, P., Tyson, J.A., & Rhee, G., 1997, *ApJ*, 583, L79
- Borgeest, U., & Refsdal, S., 1984, *A&A*, 141, 318
- Burud, I., 2002, PhD Thesis, Institut d’Astrophysique, Liege,
http://vela.astro.ulg.ac.be/themes/dataproc/deconv/theses/burud/ibthese_e.html
- Burud, I., Courbin, F., Magain, P., et al., 2002, *A&A*, 383, 71
- Burud, I., Hjorth, J., Jaunsen, A.O., et al., 2000, *ApJ*, 544, 117
- Chartas, G., Gupta, V., Garmire, G., Jones, C., Falco, E.E., Shapiro, I.I., & Tavecchio, F., 2002, *ApJ*, 565, 96
- Courbin, F., Meylan, G., Kneib, J.P., & Lidman, C., 2002, *ApJ* in press [astro-ph/0202026]
- Fabbiano, G. 1989, *ARA&A*, 27, 87
- Falco, E.E., Kochanek, C.S., Lehar, J., McLeod, B.A., Munoz, J.A., Impey, C.D., Keeton, C.R., Peng, C.Y., & Rix, H.-W., 2001, *Gravitational Lensing: Recent Progress and Future Goals*, ASP237, T. Brainerd & C.S. Kochanek, eds., (ASP: Provo) 25
- Falco, E.E., Gorenstein, M.V., & Shapiro, I.I., 1985, *ApJ*, 289, L1
- Faure, C., Courbin, F., Kneib, J.P., Alloin, D., Bolzonella, M., & Burud, I., 2002, *A&A* in press [astro-ph/0201057]
- Freedman, W.L., Madore, B.F., Gibson, B.K., et al., 2001, *ApJ*, 553, 47

- Gerhard, O., Kronawitter, A., Saglia, R.P., & Bender, R., 2001, *AJ*, 121, 1936
- Gorenstein, M.V., Falco, E.E., & Shapiro, I.I., 1988, *ApJ*, 327, 693
- Guzik, J., & Seljak, U., 2002, *MNRAS* submitted [astro-ph/0201448]
- Hjorth, J., Burud, I., Jaunsen, A.O., Schechter, P.L., Kneib, J.-P., Anderson, M.I., Korhonen, H., Clasen, J.W., Kaas, A.A., Ostensen, R., Pelt, J., & Pijpers, F.P., 2002, *ApJL* in press [astro-ph/0205124]
- Impey, C.D., Falco, E.E., Kochanek, C.S., Lehar, J., McLeod, B.A., Rix, H.-W., Peng, C.Y., & Keeton, C.R., 1998, *ApJ*, 509, 551
- Keeton, C.R., & Kochanek, C.S., 1997, *ApJ*, 487, 42
- Keeton, C.R., Kochanek, C.S., Seljak, U., 1997, *ApJ*, 482, 604
- Keeton, C.R., Falco, E.E., Impey, C.D., Kochanek, C.S., Lehar, J., McLeod, B.A., Rix, H.-W., Munoz, J.A., & Peng, C.Y., 2000, *ApJ*, 542, 74
- Kochanek, C.S., 1991, *ApJ*, 373, 354
- Kochanek, C.S., 1997, *ApJ*, 491, 13
- Kochanek, C.S., 2002, *ApJ* submitted [astro-ph/0204043]
- Kochanek, C.S., 2002, in preparation
- Koopmans, L.V.E., de Bruyn, A.G., & Jackson, N., 1998, 295, 534
- Koopmans, L.V.E., & Fassnacht, C.D., 1999, *ApJ*, 527, 513
- Koopmans, L.V.E., de Bruyn, A.G., Xanthopoulos, E., & Fassnacht, C.D., 2000, *A&A*, 356, 391
- Kundic, T., Turner, E.L., Colley, W.N., Gott, J.R., Rhoads, J.E., Wang, Y., Bergeron, L.E., Gloria, K.A., Long, D.C., Malhotra, S., & Wambsganss, J., 1997, *ApJ*, 482, 75
- Lehar, J., Falco, E.E., Kochanek, C.S., McLeod, B.A., Munoz, J.A., Impey, C.D., Rix, H.-W., Keeton, C.R., & Peng, C.Y., 2000, *ApJ*, 536, 584
- Lovell, J.E.J., et al., 1998, *ApJ*, 508, 51
- Lowenstein, M., & White, R.E., 1999, *ApJ*, 518, L50
- Maller, A.H., Simard, L., Guhathakurta, P., Hjorth, J., Jaunsen, A.O., Flores, R.A., & Primack, J.R., 2000, *ApJ*, 533, 194
- Morgan, N.D., Chartas, G., Malm, M., Bautz, M.W., Burud, I., Hjorth, J., Jones, S.E., & Schechter, P.L., 2001, *ApJ*, 555, 1

- Munoz, J.A., Kochanek, C.S., & Keeton, C.R., 2001, *ApJ*, 558, 657
- Oguri, M., Taruya, A., Suto, Y., & Turner, E.L., 2002, *ApJ*, 568, 488
- Refsdal, S., 1964, *MNRAS*, 128, 307
- Refsdal, S., & Surdej, J., 1994, *Rep. Prog. Phys.*, 57, 117
- Rix, H.-W., de Zeeuw, P.T., Cretton, N., van der Marel, R.P., & Carollo, C.M., 1997, *ApJ*, 488, 702
- Romanowsky, A.J., & Kochanek, C.S., 1999, *ApJ*, 516, 18
- Rusin, D., 2000 [*astro-ph/0004241*]
- Saha, P., 2000, *AJ*, 120, 1654
- Schechter, P.L., et al., 1997, *ApJ*, 475, L85
- Schechter, P.L., 2000, *IAU 201*, A.N. Lasenby & A. Wilkinson, eds. [*astro-ph/0009048*]
- Schild, R., & Thomson, D.J., 1997, *AJ*, 113, 130
- Schneider, P., Ehlers, J., & Falco, E.E., 1992, *Gravitational Lenses* (Berlin: Springer-Verlag)
- Treu, T., & Koopmans, L., 2002, *ApJ* submitted [*astro-ph/0202342*]
- Trotter, C.S., Winn, J.N., & Hewitt, J.N., 2000, *ApJ*, 535, 671
- Williams, L.L.R., & Saha, P., 2000, *AJ*, 119, 439
- Winn, J.N., Kochanek, C.S., McLeod, B.A., Falco, E.E., Impey, C.D., Rix, H.-W., 2002, *ApJ* in press [*astro-ph/0201551*]
- Witt, H.J., Mao, S., & Schechter, P.L., 1995, *ApJ*, 443, 18
- Witt, H.J., Mao, S., & Keeton, C.R., 2000, *ApJ*, 544, 98
- Wucknitz, O., 2002, *MNRAS* in press [*astro-ph/0202376*]
- Zhao, H.S. & Pronk, D., 2001, *MNRAS*, 320, 401